




British Thyroid Association



The Diagnosis and Management of Primary Hypothyroidism

A statement made on behalf of

The Royal College of Physicians
in particular its Patient and Carer Network
and the Joint Specialty Committee
for Endocrinology & Diabetes

The Association for Clinical Biochemistry

The Society for Endocrinology

The British Thyroid Association

The British Thyroid Foundation Patient Support Group

**The British Society of Paediatric Endocrinology
and Diabetes**

Endorsed by
the Royal College of General Practitioners

Hypothyroidism, underactivity of the thyroid gland, is common. It can make people unwell and should be treated with thyroxine (T4) tablets. Symptoms of hypothyroidism, for example tiredness, are not specific for thyroid underactivity and occur in many other situations. It is important to diagnose hypothyroidism with a blood test, because it can be dangerous to take T4 or other thyroid hormones if they are not needed. We are therefore very concerned that some patients with and without thyroid disease are being inappropriately diagnosed and managed, using thyroxine and other thyroid hormones, in ways which compromise patient safety. This is potentially an enormous problem, given that in any one year one in four of the population has their thyroid function checked.

The vast majority of patients with suspected thyroid disease are supported very well in primary care by their General Practitioners and their condition, hypothyroidism or otherwise, is appropriately diagnosed and well managed. However some patients are inappropriately diagnosed as being hypothyroid (often outside the NHS) and are started on thyroxine or other thyroid hormones which will not only cause them possible harm but leaves the true cause of their symptoms undiagnosed and therefore untreated.

This statement refers only to primary hypothyroidism. Secondary hypothyroidism is a different condition and should be managed by accredited endocrinologists in the same way as all other pituitary diseases.

DIAGNOSIS OF PRIMARY HYPOTHYROIDISM

- (a) The symptoms of hypothyroidism are very common, both in many other conditions and even in states of normal health. It is therefore essential that thyroid function is tested biochemically alongside a careful clinical assessment of the individual patient. Clinical symptoms and/or signs alone are insufficient to make a diagnosis of hypothyroidism.
- (b) The only validated method of testing thyroid function is on blood, which must include serum TSH and a measure of free thyroxine (T4).
- (c) There is no evidence to support the use of thyroid hormone testing in urine, saliva, etc or the measurement of basal body temperature in the diagnosis of thyroid dysfunction.
- (d) The results of blood tests for thyroid function can be influenced by other factors, for example in some illnesses which do not permanently damage the thyroid gland. In this case the tests will return to normal after the illness and thyroid hormone therapy is not needed (and can be harmful).
- (e) We recognise that different test methods can give different results and we support the international initiative for greater harmonisation of reference ranges and of the units used in expressing results.

TREATMENT OF PRIMARY HYPOTHYROIDISM

- (a) The aim of the treatment of hypothyroidism is to render the patient back to the normal or 'euthyroid' state.
- (b) When a sufficient dose of thyroid treatment is given to lower the TSH to the normal range (reference range) for the test method used, patients usually lose their symptoms of hypothyroidism.
- (c) Fine-tuning of TSH levels inside the reference range may be needed for individual patients.
- (d) Patients with continuing symptoms after appropriate thyroxine treatment should be further investigated to diagnose and treat the cause.
- (e) Overwhelming evidence supports the use of Thyroxine (T4) alone in the treatment of hypothyroidism. Thyroxine is usually prescribed as levothyroxine. We do not recommend the prescribing of additional Tri-iodothyronine (T3) in any presently available formulation, including Armour thyroid, as it is inconsistent with normal physiology, has not been scientifically proven to be of any benefit to patients, and may be harmful.
- (f) There are potential risks from T3 therapy, using current preparations, on bone (eg osteoporosis) and the heart (eg arrhythmia). We note that the extract marketed as Armour thyroid contains an excessive amount of T3 in relation to T4. Over-treatment with T4, when given alone, has similar risks.

TREATMENT OF SUB-CLINICAL HYPOTHYROIDISM

- (a) Sub-clinical hypothyroidism is defined as being present in a patient when the TSH is above the upper limit of the reference range but Free T4 levels are within the reference range.
- (b) Some patients, particularly those whose TSH level is greater than 10mU/l, may benefit from treatment with thyroxine in the same way as for hypothyroidism as above, as indicated in national guidelines (Thyroid function testing, Association of Clinical Biochemists, British Thyroid Association, British Thyroid Foundation, July 2006: http://www.british-thyroid-association.org/TFT_guideline_final_version_July_2006.pdf).

PATIENTS WITH NORMAL THYROID FUNCTION TESTS

- (a) We recommend that those patients whose thyroid blood tests are within the reference ranges but who have continuing symptoms, whether on thyroxine or not, should be further investigated for the non-thyroid cause of the symptoms.
- (b) A further opinion or help with these patients may be sought from appropriate specialists on specialist registers of the Royal College of Physicians or the Royal College of Paediatrics and Child Health.

CONCLUSION

- (a) Patients with suspected primary hypothyroidism should only be diagnosed with blood tests including measurement of TSH.
- (b) Patients with primary hypothyroidism should be treated with T4 using levothyroxine tablets (BNF) alone.
- (c) There is no indication for the prescription of T4 or any preparation containing thyroid hormones to patients with thyroid blood tests within the reference ranges.
- (d) In patients with suspected primary hypothyroidism there is no indication for the prescription of T4 or any preparation containing thyroid hormones to patients with thyroid blood tests initially within the normal range. Thus patients with normal T4 and TSH do not have primary hypothyroidism and even if they have symptoms which might suggest this should not be given thyroid hormone replacement therapy.
- (e) The College does not support the use of thyroid extracts or thyroxine and T3 combinations without further validated research published in peer-reviewed journals. Therefore, the inclusion of T3 in the treatment of hypothyroidism should be reserved for use by accredited endocrinologists in individual patients.
- (g) Laboratories which measure thyroid function in other bodily fluids besides blood need to provide analytical and clinical validation to demonstrate their efficacy.
- (h) The above statements reflect best practice of clinical endocrinologists accredited by the Royal College of Physicians and the Royal College of Paediatrics and Child Health.

19th November 2008

Copyright

All rights reserved. No part of this publication may be reproduced in any form (including photocopying or storing it in any medium by electronic means and whether or not transiently or incidentally to some other use of this publication) without the written permission of the copyright owner. Applications for the copyright owner's permission to reproduce any part of this publication should be addressed to the publisher.

Copyright © 2009 Royal College of Physicians 11 St Andrews Place, London NW1 4LE

www.rcplondon.ac.uk

Registered Charity No 210508